

ON THE MIN-PROJECTIVE MODULES

M. Amini, A. Farajzadeh and S. Bayati

Received: 10 March 2010; Revised: 06 April 2011

Communicated by Abdullah Harmanci

ABSTRACT. Let R be a commutative ring. An R -module M is called min-projective if $\text{Ext}_R^1(M, \frac{R}{I}) = 0$, for every simple ideal I . In this paper, we first give some results of min-projective R -modules on the some specific rings such as cotorsion rings, von Neumann regular rings and coherent rings. Then we investigate min-projective covers on universally min-projective rings. Finally, we deal with some characterizations of min-projective modules over a perfect ring.

Mathematics Subject Classification (2010): 16D50, 16E50

Keywords: min-projective modules, min-flat modules, universally min-projective rings

1. Introduction

Throughout this paper, R denotes a commutative ring. Let \mathcal{C} be a class of R -modules and G be an element of \mathcal{C} . The homomorphism $\theta : G \rightarrow K$ with $G \in \mathcal{C}$ is called a \mathcal{C} -precover of K if for any homomorphism $f : G' \rightarrow K$ with $G' \in \mathcal{C}$, there exists a homomorphism $g : G' \rightarrow G$ such that $\theta \circ g = f$. Moreover, if the mapping g is automorphism of G when $G = G'$ and $f = g$ then \mathcal{C} -precover θ is called the \mathcal{C} -cover of G . One can similarly define the dual of the \mathcal{C} -preenvelope and \mathcal{C} -envelope, see [6], for more details. An R -module C is said to be cotorsion if $\text{Ext}_R^1(F, C) = 0$, for all flat R -module F , see [15, Definition 3.1.1]. There are many papers which deal with to the concept of projective modules and injective modules and the related topics, see for instance [2,4,10,11]. An R -module N is called *finitely presented* if there exists the exact sequence $R^{(n)} \rightarrow R^{(m)} \rightarrow N \rightarrow 0$ and an R -module M is called *FP-injective* if $\text{Ext}_R^1(N, M) = 0$ for any finitely presented R -module N . Also, an R -module N is called *FP-projective* if $\text{Ext}_R^1(N, M) = 0$ for any *FP-injective* R -module M . We refer the reader to [4] for more details about *FP-projective* and *FP-injective* modules. Note that for any R -module M , $\sigma_M : M \rightarrow C(M)$, $\tau_M : M \rightarrow FE(M)$ and $\xi_M : F(M) \rightarrow M$ will denote, respectively, a cotorsion envelope, an *FP-injective* preenvelope and a flat cover for

M . An R -module M is called *min-projective* if $\text{Ext}_R^1(M, \frac{R}{I}) = 0$ for any simple ideal I . An R -module N is called *min-flat* if $\text{Tor}_1^R(N, \frac{R}{I}) = 0$ for any simple ideal I and if every R -module is min-projective or min-flat, then R is called a *universally min-projective* ring or a *universally min-flat* ring. Recall that a ring R is called *coherent* if every finitely generated ideal is finitely presented, see [15, Definition 1.1.4]. The *socle* of R , denoted by $\text{Soc}(R)$, is the direct sum of nonzero simple ideals of R . We use $\text{Soc}(R) \leq_e R$ to mean that $\text{Soc}(R)$ is an essential ideal of R and $r \in R$ is singular if $\text{Ann}_R(r) \leq_e R$. A ring R is called *von Neumann regular* if for each $r \in R$, there is $r' \in R$ with $rr'r = r$, see [12]. A ring R is said to be *perfect* when every R -module has a projective cover, see [15]. In this paper, some characterizations of min-projective modules on cotorsion rings, von Neumann regular rings, coherent rings, universally min-projective rings and perfect rings are given. For instance, it is shown that R is a cotorsion ring if and only if every flat R -module is min-projective; R is a von Neumann regular ring if and only if R is a coherent ring and every *FP*-projective R -module is min-projective; on universally min-flat rings with $\text{Soc}(R) \leq_e R$, R is a universally min-projective ring if and only if every min-flat R -module has an Ω -cover with the unique mapping property if and only if R is a cotorsion ring with $Z(R) = 0$, where Ω is the class of min-projective R -modules and $Z(R)$ is the set of all singular elements. Also, we prove that R is a perfect ring if and only if every min-projective R -module is cotorsion if and only if every flat R -module is min-projective and every min-projective R -module has a cotorsion envelope with the unique mapping property if and only if for each R -homomorphism $f : M_1 \rightarrow M_2$ with M_1 and M_2 min-projective, $\ker(f)$ is cotorsion.

2. Main Results

We start by the following definition.

Definition 2.1. Let R be a ring. An R -module M is called *min-projective* if $\text{Ext}_R^1(M, \frac{R}{I}) = 0$ for any simple ideal I .

It is well-known that if R is a cotorsion R -module, then R is a cotorsion ring. The following proposition shows that, on cotorsion rings, every flat module is a min-projective module.

Proposition 2.2. *Let R be a ring. Then the following statements are equivalent:*

- (1) R is a cotorsion ring;
- (2) Every flat R -module is a min-projective.

Proof. (1) \Rightarrow (2) It is known from [5, Lemma 2.14], which I is a cotorsion ring. Now consider the exact sequence $0 \rightarrow I \rightarrow R \rightarrow \frac{R}{I} \rightarrow 0$. Then for every flat R -module M , we get the exact sequence $0 = \text{Ext}_R^1(M, R) \rightarrow \text{Ext}_R^1(M, \frac{R}{I}) \rightarrow \text{Ext}_R^2(M, I) = 0$. Hence $\text{Ext}_R^1(M, \frac{R}{I}) = 0$ and so M is min-projective R -module. (2) \Rightarrow (1) is clear. \square

It is obvious that every projective module is a min-projective module. However, the following example shows that the converse is not true in general. Before this, we recall that a ring is said to be *hereditary* if all of its ideals are projective, see [12]. If R has no simple ideal, then the socle of R is defined to be zero and in this case it is clear that every R -module is a min-projective module. The following example shows that the definition of min-projective R -modules is a proper generalization of projective modules.

Example 2.3. *Let R be a ring.*

- (a) *If R is a non-hereditary ring such that $\text{Soc}(R) = (0)$, then some of ideals of R are min-projective, while they are not projective R -modules. In particular, if $R \cong \frac{K[x_n : n \geq 1]}{(x_i x_j : i \geq 1 \text{ and } j \geq 1)}$, then $\text{Soc}(R) = (0)$ and so for every $n \geq 1$, the ideal (x_n) is a min-projective module but it is not projective.*
- (b) *Let R be a reduced ring, that is R has no non-zero nilpotent element which is not decomposable (for example R can be an integral domain which is not a field). We show that R contains no simple ideal. By contrary, suppose that R contains a simple ideal, say Re . Then since R is reduced, we deduce that e is an idempotent element and so by Brauer's lemma (see [8, 10.22]), R is decomposable that is a contradiction. Hence every R -module is min-projective, because R contains no simple ideal. But not all R -modules are projective.*
- (c) *Let $R \cong D_1 \times \cdots \times D_n$, where every D_i , $1 \leq i \leq n$, is an integral domain which is not field. Then every R -module is min-injective. But there are R -modules which are not projective.*
- (d) *From Part (c), we conclude that any module over the ring \mathbb{Z} of integers is min-projective. But not all \mathbb{Z} -modules are projective.*

In the following proposition, some properties of modules on cotorsion ring are studied.

It is trivial that min-projective modules are closed under extensions over any ring. So, we have the following proposition.

Proposition 2.4. *Let R be a cotorsion ring. Then*

- (1) *Let $\alpha : N \rightarrow M$ be a monomorphism. Then $\text{coker}(\alpha)$ is min-projective if and only if $\text{coker}(\sigma_M \alpha)$ is min-projective.*
- (2) *Let N be a submodule of M . If M is min-projective and $\frac{M}{N}$ is flat, then N is also min-projective.*
- (3) *Every cotorsion envelope of a min-projective R -module is min-projective.*

Proof. (1) and (3) are clear, by the fact which was mentioned before the proposition. Now, we prove (2). Let I be a simple ideal. Then the short exact sequence $0 \rightarrow N \rightarrow M \rightarrow \frac{M}{N} \rightarrow 0$ induces the exact sequence

$$0 = \text{Ext}_R^1(M, \frac{R}{I}) \rightarrow \text{Ext}_R^1(N, \frac{R}{I}) \rightarrow \text{Ext}_R^2(\frac{M}{N}, \frac{R}{I}) = 0.$$

The first equality follows by Proposition 2.2. Hence $\text{Ext}_R^1(N, \frac{R}{I}) = 0$ and so N is min-projective. \square

Proposition 2.5. *A min-flat R -module M is min-projective if and only if the $\frac{R}{I}$ -module $\frac{M}{MI}$ is min-projective, for every simple ideal I .*

Proof. This follows from the isomorphism $\text{Ext}_R^1(M, \frac{R}{I}) \simeq \text{Ext}_{\frac{R}{I}}^1(\frac{M}{MI}, \frac{R}{I})$, see [13, Lemma 5.1]. \square

In the following proposition, we give some conditions under which the direct sum of a family of min-flat R -modules is min-projective. Before this, we recall that for any R -module M , the R -module $\text{Hom}_Z(M, \frac{Q}{Z})$ is denoted by M^+ .

Proposition 2.6. *Let $\{\frac{M_i}{M_i I} : i \in I\}$ and $\{\frac{M_i^{++}}{M_i^{++} I} : i \in I\}$ be two indexed sets of min-projective $\frac{R}{I}$ -modules, where I is a simple ideal. If every M_i is min-flat, then*

- (1) $\coprod_{i \in I} M_i$ is min-projective.
- (2) $\coprod_{i \in I} M_i^{++}$ is min-projective.

Proof. (1) By Proposition 2.5, we have $\text{Ext}_R^1(M_i, \frac{R}{I}) = 0$. Thus by [12, Theorem 7.13], $\text{Ext}_R^1(\coprod_{i \in I} M_i, \frac{R}{I}) \simeq \prod_{i \in I} \text{Ext}_R^1(M_i, \frac{R}{I}) = 0$ and so $\coprod_{i \in I} M_i$ is min-projective.

(2) This follows from [3, Lemma 3.2]. \square

Remark 2.7. *Let R be a coherent ring. By [6, Theorem 7.4.1], every R -module has a special FP-injective pre-envelope, i.e; there is an exact sequence $0 \rightarrow M \rightarrow F \rightarrow L \rightarrow 0$, where F is FP-injective and L is FP-projective and every R -module has a special FP-projective precover, i.e; there is an exact sequence $0 \rightarrow K \rightarrow P \rightarrow M \rightarrow 0$, where P is FP-projective and K is FP-injective.*

It is well-known that R is a von Neumann regular ring if and only if every R -module is flat, see [12, Theorem 4.9]. In the following theorem, we give a characterization of a von Neumann regular ring.

Recall that a \mathcal{C} -cover $\phi : M \rightarrow N$ has the *unique mapping property* if for any homomorphism $f : A \rightarrow N$ with $A \in \mathcal{C}$, there exists a unique $g : A \rightarrow M$ such that $\phi g = f$. One can similarly define the dual of the \mathcal{C} -envelope, see [6].

Theorem 2.8. *Let R be a ring. Then the following statements are equivalent:*

- (1) R is a von Neumann regular ring;
- (2) R is a coherent ring and every FP -projective R -module is min-projective.

Proof. (1) \Rightarrow (2) By [2, Corollary 4.3], R is a coherent ring. Let N be an R -module. Then by Remark 2.7, there is an exact sequence $0 \rightarrow K \rightarrow P \rightarrow N \rightarrow 0$, where P is FP -projective and K is FP -injective. Therefore, for every FP -projective R -module M , we obtain the exact sequence

$$0 = \text{Ext}_R^1(M, P) \rightarrow \text{Ext}_R^1(M, N) \rightarrow \text{Ext}_R^2(M, K) = 0.$$

Thus $\text{Ext}_R^1(M, N) = 0$ and so every FP -projective R -module is min-projective.

(2) \Rightarrow (1) By [15, Theorem 2.3.1], every R -module has a FP -injective envelope and by Remark 2.7 and (2), its cokernel is FP -projective. Let M be a min-projective R -module. Then there exists a commutative diagram with the exact rows:

$$\begin{array}{ccccccc} & & & & 0 & & \\ & & & & \downarrow & & \\ 0 & \longrightarrow & M & \xrightarrow{\tau_M} & FE(M) & \xrightarrow{\gamma} & L \longrightarrow 0 \\ & & & & 0 \searrow & \downarrow \tau_L \gamma & \swarrow \tau_L \\ & & & & & FE(L) & \end{array}$$

Note that by [12, Proposition 7.24] and Remark 2.7, for every FP -injective R -module N , there exists a split exact sequence $0 \rightarrow N \rightarrow D \rightarrow C \rightarrow 0$, where D is an FP -injective R -module and C is an FP -projective R -module. So for every R -module K , $\text{Ext}_R^1(K, N) = 0$. Therefore by [5, Corollary 2.12] and [12, Theorem 3.56], every finitely presented R -module is projective. Hence every R -module is FP -injective. So the FP -injective envelopes τ_L and τ_M satisfy unique mapping property. Since $\tau_L \gamma \tau_M = 0 = 0 \tau_M$, from (2), we have $\tau_L \gamma = 0$. Thus $L = \text{im}(\gamma) \subseteq \ker(\tau_L) = 0$ and hence $L = 0$. Therefore, $M = FE(M)$ and so every min-projective R -module is FP -injective. Thus (1) follows from (2) and [2, Corollary 4.3] and the proof completes. \square

Definition 2.9. Let R be a ring. A ring R is called *universally min-projective* if every R -module is min-projective.

Definition 2.10. Let R be a ring. A ring R is called *universally min-flat* when every R -module is min-flat.

Now, we present the following characterizations of the universally min-projective rings. From now and for simplicity, we denote the class of min-projective R -modules by Ω .

Theorem 2.11. *Let R be a ring. Then the following statements are equivalent:*

- (1) *For any simple ideal I and any flat R -module M , the $\frac{R}{I}$ -module $\frac{M}{MI}$ is min-projective;*
- (2) *R is a cotorsion ring;*
Moreover, if R is a universally min-flat ring with $\text{Soc}(R) \leq_e R$, then the above conditions are equivalent to:
- (3) *R is a cotorsion ring with $Z(R) = 0$;*
- (4) *For every simple ideal I of R , $\frac{R}{I}$ is cotorsion and $J(R) = 0$;*
- (5) *R is a universally min-projective ring;*
- (6) *Every min-flat R -module has an Ω -cover with the unique mapping property;*
- (7) *Every min-flat R -module is min-projective.*

Proof. (1) \Rightarrow (2) We shall show that $\text{Ext}_R^1(M, R) = 0$, for every flat R -module M . Let I be a simple ideal of R . Then we have the short exact sequence $0 \rightarrow I \rightarrow R \rightarrow \frac{R}{I} \rightarrow 0$. By [5, Lemma 2.14], $\text{Ext}_R^1(M, I) = 0$. Since every flat module is min-flat, Proposition 2.5 implies that M is min-projective. Hence $\text{Ext}_R^1(M, \frac{R}{I}) = 0$. Therefore, the above exact sequence induces the exact sequence

$$0 = \text{Ext}_R^1(M, I) \rightarrow \text{Ext}_R^1(M, R) \rightarrow \text{Ext}_R^1(M, \frac{R}{I}) = 0,$$

for every flat R -module M . Thus $\text{Ext}_R^1(M, R) = 0$, as desired.

(2) \Rightarrow (1) This follows from Propositions 2.2 and 2.5.

(2) \Rightarrow (3) R is a cotorsion ring by (2). Note that $\frac{R}{I}$ is flat and so by [5, Theorem 2.16] R is a *PS* ring (every simple ideal of R is projective). Therefore, $\text{Soc}(R)$ is projective and hence by [9, Exercise 12 (A), p.269], $Z(\text{Soc}(R)) = 0$. Therefore, by [1, Lemma 7.2], $Z(\text{Soc}(R)) = Z(R) \cap \text{Soc}(R) = 0$ and so $Z(R) = 0$.

(3) \Rightarrow (4) Let I be a simple ideal of R and M be a flat R -module. Then we obtain the exact sequence $0 \rightarrow I \rightarrow R \rightarrow \frac{R}{I} \rightarrow 0$ which gives rise to the exactness of

$$\cdots \rightarrow \text{Ext}_R^1(M, R) \rightarrow \text{Ext}_R^1(M, \frac{R}{I}) \rightarrow \text{Ext}_R^2(M, I) \rightarrow \cdots$$

By (3), R is cotorsion. So, $Ext_R^1(M, R) = 0$. Also, from [5, Lemma 2.14], we conclude that $Ext_R^2(M, I) = 0$. Therefore, $Ext_R^1(M, \frac{R}{I}) = 0$ and so $\frac{R}{I}$ is cotorsion. Now, we claim that $J(R) = Ann_R(Soc(R)) = Z(R)$. Since the annihilator of any simple ideal is a maximal ideal, we deduce that $J(R) = Ann_R(Soc(R))$. It is clear that $Soc(R)^2 = 0$. Thus $Soc(R) \subseteq Ann(Soc(R))$. Now, since $Soc(R) \leq_e R$, we deduce that $Ann_R(Soc(R)) \leq_e R$ and so $Ann_R(Soc(R)) \subseteq Z(R)$. Since R is a cotorsion ring, we deduce that $\frac{R}{J(R)}$ is semisimple, by [7, Theorem 6]. Thus $Z(\frac{R}{J(R)}) = 0$ and hence $Z(R) \subseteq J(R)$. So, by (3), $J(R) = 0$.

(4) \Rightarrow (5) Note that R is a von Neumann regular ring by [5, Theorem 2.16] and so by [5, Corollary 2.12], $\frac{R}{I}$ is injective. Hence (5) follows.

(5) \Rightarrow (6) This is clear.

(6) \Rightarrow (7) Let M be a min-flat R -module. Then there is a commutative diagram with exact rows:

$$\begin{array}{ccccccc} & & & N' & & & \\ & & & \phi \swarrow \alpha\phi \downarrow & \searrow & & 0 \\ 0 & \longrightarrow & K & \xrightarrow{\alpha} & N & \xrightarrow{\psi} & M \longrightarrow 0, \\ & & & \downarrow & & & \\ & & & 0 & & & \end{array}$$

where ψ and ϕ are Ω -cover with the unique mapping property. Since $\psi\alpha\phi = 0 = \psi$, we have $\alpha\phi = 0$ by (6). Therefore, $K = im(\phi) \subseteq ker(\alpha) = 0$ and so $K = 0$. Thus $M = N$ and hence every min-flat R -module is min-projective.

(7) \Rightarrow (1) Let I be a simple ideal of R and M be a flat R -module. Then it is clear that M is a min-flat R -module. So, by (7), M is a min-projective R -module. Thus Proposition 2.5 implies that $\frac{M}{MI}$ is min-projective $\frac{R}{I}$ -module and so we are done. \square

Corollary 2.12. *Let R be a coherent ring. Then the following statements are equivalent:*

- (1) R is a universally min-projective ring;
- (2) Every R -module has an Ω -cover with the unique mapping property;
- (3) For every simple ideal I , $\frac{R}{I}$ is cotorsion and every FP -projective R -module is min-projective.

Proof. (1) \Rightarrow (2) It is trivial.

(2) \Rightarrow (3) By (2) and (7) of Theorem 2.11, $\frac{R}{I}$ is cotorsion. Let M be a FP -projective R -module, similar to proof (6) \Rightarrow (7) of Theorem 2.11, R -module M is min-projective and so (3) follows.

(3) \Rightarrow (1) This follows from Theorem 2.8 and [5, Corollary 2.12]. \square

Corollary 2.13. *Let R be a coherent ring such that $\frac{R}{I}$ be an injective R -module, for every simple ideal of R . If $h : M \rightarrow N$ is a homomorphism of min-projective R -modules, $\text{coker}(h)$ is min-projective.*

Proof. By Theorem 2.8 and Corollary 2.12, M is FP -injective. So, the short exact sequence $0 \rightarrow M \xrightarrow{h} N \rightarrow \frac{N}{\text{im}(h)} \rightarrow 0$ is pure. Therefore, for every R -module B , $\text{Tor}_1^R(B, \frac{N}{\text{im}(h)}) = 0$ and so by Proposition 2.2, $\frac{N}{\text{im}(h)}$ is min-projective. \square

Corollary 2.14. *Let R be a coherent ring and $\frac{R}{I}$ be a cotorsion module, for every simple ideal I . Then R is a von Neumann regular ring if and only if R is a universally min-projective ring.*

Proof. This is a direct consequence of Theorem 2.8 and Corollary 2.12. \square

From [14, Proposition 9.43], we know that R is a perfect ring if and only if every flat R -module is projective. In the following theorem, we give some other characterizations of perfect rings.

Theorem 2.15. *Let R be a ring. Then the following statements are equivalent:*

- (1) R is a perfect ring;
- (2) Every min-projective R -module is cotorsion;
- (3) Every flat R -module is min-projective and every min-projective R -module has a cotorsion envelope with the unique mapping property;
- (4) For each R -homomorphism $f : M_1 \rightarrow M_2$ with M_1 and M_2 min-projective, $\ker(f)$ is cotorsion;
- (5) For each min-projective R -module M , the functor $\text{Hom}_R(-, M)$ is exact with respect to each pure exact sequence $0 \rightarrow K \rightarrow P \rightarrow L \rightarrow 0$ in which P is projective. In addition, L and $\frac{R}{I}$ -module $\frac{K}{KI}$ are min-projective, for every simple ideal I .

Proof. (1) \Rightarrow (2) For every flat R -module F and every min-projective R -module M , $\text{Ext}_R^1(F, M) = 0$. So (2) follows.

(2) \Rightarrow (1) In the short exact sequence $0 \rightarrow K \rightarrow P \xrightarrow{\xi_L} L \rightarrow 0$ with L flat, P projective and ξ_L projective cover of L , K is cotorsion. Thus (2) implies that P is cotorsion. So, for every flat R -module F , we obtain the exact sequence

$$0 = \text{Ext}_R^1(F, P) \rightarrow \text{Ext}_R^1(F, L) \rightarrow \text{Ext}_R^2(F, K) = 0.$$

Hence $\text{Ext}_R^1(F, L) = 0$ and so every flat R -module is cotorsion and by [15, Proposition 3.3.1], (1) follows.

(1) \Rightarrow (3) This is a direct consequence of [14, Proposition 9.43] and [5, Theorem 2.18].

(3) \Rightarrow (1) Let $C(M)$ be the cotorsion envelope of min-projective R -module M . There is the following commutative diagram:

$$\begin{array}{ccccccc} & & & & 0 & & \\ & & & & \downarrow & & \\ 0 & \longrightarrow & M & \xrightarrow{\sigma_M} & C(M) & \xrightarrow{\gamma} & L \longrightarrow 0 \\ & & & & \searrow \sigma_L \gamma & & \swarrow \sigma_L \\ & & & & C(L) & & \end{array}$$

Note that by [15, Theorem 3.4.2], L is flat and so σ_L exists. Hence $\sigma_L \gamma \sigma_M = 0 = 0 \sigma_M$ and so $\sigma_L \gamma = 0$. Therefore, $L = \text{im}(\gamma) \subseteq \ker(\sigma_L) = 0$ and so $M = C(M)$. Thus by (2), R is a perfect ring.

(1) \Rightarrow (4) This follows from the fact that every module is cotorsion over perfect rings.

(4) \Rightarrow (1) Let M be a min-projective R -module. From the above commutative diagram, we have $M = \ker(\gamma) = \ker(\sigma_L \gamma)$. Thus by (4), M is cotorsion. So, (1) follows from (2).

(1) \Rightarrow (5) For every R -module B , we obtain the exact sequence

$$0 \longrightarrow B \otimes_R K \longrightarrow B \otimes_R P \longrightarrow B \otimes_R L \longrightarrow 0.$$

Hence $\text{Tor}_1^R(L, B) = 0$ and so L is flat. Then for any min-projective R -module M , we have the following exact sequence

$$\text{Hom}_R(P, M) \longrightarrow \text{Hom}_R(K, M) \longrightarrow \text{Ext}_R^1(L, M) = 0.$$

By (2), M is cotorsion; therefore, the functor $\text{Hom}_R(-, M)$ is exact. By [14, Proposition 9.43], L is min-projective. Thus for every simple ideal I , we get the exact sequence

$$0 = \text{Ext}_R^1(P, \frac{R}{I}) \longrightarrow \text{Ext}_R^1(K, \frac{R}{I}) \longrightarrow \text{Ext}_R^2(L, \frac{R}{I}) = 0.$$

Hence $\text{Ext}_R^1(K, \frac{R}{I}) = 0$ and hence K is min-projective. Since P and L are flat, we deduce that K is flat and subsequently by Proposition 2.5, $\frac{K}{KI}$ is min-projective.

(5) \Rightarrow (1) For every min-projective R -module M and every flat R -module L , we have $\text{Ext}_R^1(L, M) = 0$. So, every min-projective R -module is cotorsion. Thus (1) follows from (2). \square

Example 2.16. Let \mathbb{Z} be the ring of integer numbers. Then we show that the \mathbb{Z} -module \mathbb{Z} is min-projective which is not cotorsion. Suppose to the contrary, \mathbb{Z}

is cotorsion. Then by Proposition 2.2, every flat R -module is min-projective and so every flat R -module is cotorsion. Hence [15, Proposition 3.3.1] implies that \mathbb{Z} is a perfect ring and this contradicts this fact that \mathbb{Z} is not a perfect ring, see [12, Example 4.61].

Acknowledgment. The authors would like to thank the editor and the referee for their careful reading of the paper and for many remarks and suggestions which led to the improvements in the paper. Also, the authors wish to express their sincere thanks to F. Shaveisi, for the valuable suggestions and the careful reading of the paper, which improve the presentation of the article and help correct many mistakes.

References

- [1] F. W. Anderson and K. R. Fuller, *Rings and Categories of Modules*, Springer-Verlag, New York, 1974.
- [2] N. Q. Ding and L. X. Mao, *FP-projective dimensions*, *Comm. Algebra*, 33 (2005), 1153–1170.
- [3] N. Q. Ding and L. X. Mao, *Min-flat modules and min-coherent rings*, *Comm. Algebra*, 36 (2007), 635–650.
- [4] N. Q. Ding and L. X. Mao, *Notes on FP-projective modules and FP-injective modules*, *Advances in ring theory*, 151–166, World Sci.Publ., Hackensack, NJ, 2005.
- [5] N. Q. Ding and L. X. Mao, *Notes on the cotorsion modules*, *Comm. Algebra*, 33 (2005), 349–360.
- [6] E. E. Enochs and M. G. Jenda, *Relative Homological Algebra*, Walter de Gruyter, Berlin, New York, 2000.
- [7] P. A. Guil Asensio and I. Herzog, *Left cotorsion rings*, *Bull. London Math. Soc.*, 36 (2004) 303-309.
- [8] T. Y. Lam, *A First Course in Non-Commutative Rings*, Springer-Verlag, New York, 1991.
- [9] T. Y. Lam, *Lectures on Modules and Rings*, Springer-Verlag, Berlin, 1999.
- [10] M. J. Nikmehr, F. Shaveisi and R. Nikandish, *n-Projective modules*, *Algebras, Groups and Geometries*, 24 (2007), 447–454.
- [11] M. J. Nikmehr and F. Shaveisi, *T-dimension and $(n + \frac{1}{2}, T)$ -Projective modules*, *Southeast Asian Bull. Math.*, 35 (2011), 1–11.
- [12] J.J. Rotman, *An Introduction to Homological Algebra*, Academic Press, New York, 1979.

- [13] F. L. Sandomierski, *Homological dimensions under change of rings*, Math.Z., 130 (1973), 55–65.
- [14] R. Wisbauer, *Foundation of Module and Ring Theory*, Gordon and Breach Publishers, 1991.
- [15] J. Xu, *Flat covers of modules*, Lecture Notes in Mathematics, 1643, Berlin, Springer-Verlag, 1996.

M. Amini

Department of Mathematics
Faculty of Science
Payame Noor University
Sonqor, Iran
e-mail: mamini1356@yahoo.com

A. Farajzadeh

Islamic Azad university
Kermanshah Branch, Iran
e-mail: faraj1348@yahoo.com

S. Bayati

Department of Mathematics
Faculty of Science
Payame Noor University
Sonqor, Iran
e-mail: s.bayati@gmail.com